

Foundations of Computer Science

Comp109

University of Liverpool
Boris Konev
konev@liverpool.ac.uk
<http://www.csc.liv.ac.uk/~konev/COMP109>

Part 6. Combinatorics

Comp109 Foundations of Computer Science

Reading

[Discrete Mathematics and Its Applications](#), K. H. Rosen, Sections 6.1, 6.3, 6.4

Contents

- Basics of counting
- Notation for sums and products. The factorial function.
- Counting permutations and combinations.
- Binomial coefficients.

Developing ideas (1)

All chairs in a room are labelled with a single digit followed by a lower-case letter. What is the largest number of differently numbered chairs?



Developing ideas (2)

How many different bit strings of length 8 are there?

- How many different bytes are there?

0000 0000, 0000 0001, 0000 0010, 0000 0011, ...

Developing ideas (3)

How many ways there are to select 3 students for a prospectus photograph (order matters) from a group of 5?




The product rule

If there is a sequence of k events with n_1, \dots, n_k possible outcomes, then the total number of outcomes for the sequence of k events is


$$n_1 \times n_2 \times \dots \times n_k.$$

Example


How many distinct car licence plates are there consisting of six characters, the first three of which are letters and the last three of which are digits?

Developing ideas (4)	Disjoint events	The sum rule
<p>How many ways there are to select a male and a female student for a prospectus photograph (order matters) from a group of 2 male and 3 female students?</p> 	<p>Two events are said to be disjoint (or “mutually exclusive”) if they can’t occur simultaneously.</p> <p>Example: If you have 3 pairs of blue jeans and 2 pairs of black jeans, then there are $3 + 2 = 5$ different pairs of jeans which are blue or black which you could wear.</p>	<p>If A and B are disjoint events and there are n_1 possible outcomes for event A and n_2 possible outcomes for event B then there are $n_1 + n_2$ possible outcomes for the event “either A or B”.</p>

Example	Example	Set-theoretic interpretation
<p>How many three-digit numbers begin with 3 or 4?</p>	<p><i>I wish to take two pieces of fruit with me for lunch. I have three bananas, four apples and two pears. How many ways can I select two pieces of fruit of different type?</i></p>	<ul style="list-style-type: none"> ■ If A and B are disjoint sets (that is, $A \cap B = \emptyset$) then $A \cup B = A + B$. ■ Any sequence of k events can be regarded as an element of the Cartesian product $A_1 \times \dots \times A_k$. This set has size $A_1 \times \dots \times A_k$.

Developing ideas (5)	Answer	Note: lazy users
<p>A computer password is a string of 8 characters, where each character is an uppercase letter or a digit. Each password must contain at least one digit.</p> <p>How many different passwords are there?⁴</p> 	<p>2, 612, 282, 842, 880</p>	<p>How many different 8-character passwords can be obtained by combining 3-letter word, a 4-letter word and a digit? (According to http://www.scrabblefinder.com there are 1015 3-letter and 4030 4-letter English words.)</p>

Answer
<p>245,427,000 (about 0.009%)</p> <p>Beware of passwords like HOT4FUZZ</p>


Developing ideas (6)
<p>How many bit strings of length 8 either start with 1 or finish with 00?</p>  $ A \cup B = A + B - A \cap B $

The subtraction rule
<p>If there are n_1 possible outcomes for event A, n_2 possible outcomes for event B and n_3 of these outcomes are shared between A and B then there are</p> $n_1 + n_2 - n_3$ <p>possible outcomes for the event “A or B”.</p>

http://www.csc.liv.ac.uk/~konev/COMP109	Part 6. Combinatorics	18 / 44
---	-----------------------	---------

http://www.csc.liv.ac.uk/~konev/COMP109	Part 6. Combinatorics	19 / 44
---	-----------------------	---------

http://www.csc.liv.ac.uk/~konev/COMP109	Part 6. Combinatorics	20 / 44
---	-----------------------	---------

Developing ideas (7)
<p>How many ways there are to select 2 representatives from a group of 5 students?</p> 

The division rule
<p>Given n possible outcomes, if</p> <ul style="list-style-type: none"> ■ some of the n outcomes are the same ■ every group of indistinguishable outcomes contains exactly d elements <p>there are n/d different outcomes.</p>

Mini summary
<p>Counting problems can be hard</p> <p>Four decomposition rules:</p> <ul style="list-style-type: none"> ■ The product rule ■ The sum rule ■ The subtraction rule ■ The division rule <p>To move further we need some mathematical notation</p>

http://www.csc.liv.ac.uk/~konev/COMP109	Part 6. Combinatorics	21 / 44
---	-----------------------	---------

http://www.csc.liv.ac.uk/~konev/COMP109	Part 6. Combinatorics	22 / 44
---	-----------------------	---------

http://www.csc.liv.ac.uk/~konev/COMP109	Part 6. Combinatorics	23 / 44
---	-----------------------	---------

Sums and products of a sequence
<p>Given a sequence of numbers</p> $\dots, a_1, a_2, a_3, \dots, a_m, a_{m+1}, \dots, a_n, \dots$ <p>we use the notation</p> <ul style="list-style-type: none"> ■ $\sum_{i=m}^n a_i$ to represent $a_m + a_{m+1} + \dots + a_n$ and ■ $\prod_{i=m}^n a_i$ to represent $a_m \times a_{m+1} \times \dots \times a_n$

Examples
<ul style="list-style-type: none"> ■ $\sum_{i=1}^5 i = 1 + 2 + 3 + 4 + 5$ ■ $\prod_{i=0}^6 i^2 = 0^2 \times 1^2 \times 2^2 \times 3^2 \times 4^2 \times 5^2 \times 6^2 = 0.$ <p>Notice that i is just an index, so</p> $\sum_{i=1}^5 i = \sum_{j=1}^5 j = \sum_{k=1}^5 k = 1 + 2 + 3 + 4 + 5$

Special case $m = n$
$\sum_{i=m}^n = a_m, \quad \text{when } m = n$ <p>For example</p> $\sum_{i=3}^3 i = 3$

http://www.csc.liv.ac.uk/~konev/COMP109	Part 6. Combinatorics	24 / 44
---	-----------------------	---------

http://www.csc.liv.ac.uk/~konev/COMP109	Part 6. Combinatorics	25 / 44
---	-----------------------	---------

http://www.csc.liv.ac.uk/~konev/COMP109	Part 6. Combinatorics	26 / 44
---	-----------------------	---------

We can express some equalities more neatly using this notation.

- In Part 1 of the module, we proved that

$$\sum_{i=0}^n i = \frac{n(n+1)}{2}.$$

- In Part 2 of the module, we defined the Cartesian product $A_1 \times \dots \times A_k$ of k sets.

$$A_1 \times \dots \times A_k = \{(a_1, a_2, \dots, a_k) \mid a_i \in A_i\}.$$

The size of the Cartesian product is $\prod_{i=1}^k |A_i|$.

Let $f : D \rightarrow \mathbb{R}$ be a function with some domain D .

Then for $S \subseteq D$,

- $\sum_{i \in S} f(i)$ denotes the sum of $f(i)$ over all $i \in S$ and
- $\prod_{i \in S} f(i)$ denotes the product of $f(i)$ over all $i \in S$.

The product $\prod_{i=1}^n i$ comes up so often that it has a name. It is called **n factorial** and is written as **$n!$** .

Examples:

- $5! = 1 \times 2 \times 3 \times 4 \times 5 = 120$.
- $3! = 1 \times 2 \times 3 = 6$.
- $1! = 1$.
- $0! = 1$.

Permutations

k -permutations

Examples

A **permutation** of a set is just an ordering of its elements.

Example: The permutations of the set $\{1, 2, 3\}$ are

- 1, 2, 3
- 2, 1, 3
- 3, 1, 2
- 1, 3, 2
- 2, 3, 1
- 3, 2, 1

By the product rule the number of permutations of an n -element set is

$$n! = n \times (n-1) \times \dots \times 1$$

because there are n choices for the first element, then $n-1$ choices for the 2nd element, then $n-2$ choices for the 3rd element, and so on.

A selection of k distinct elements of a set, where order matters, is called a **k -permutation** of the set.

The number k -permutations of an n -element set is

$$P(n, k) = n \times (n-1) \times \dots \times (n-(k-1)) = \frac{n!}{(n-k)!}.$$

- How many ways there are to select 3 students for a prospectus photograph (order matters) from a group of 5?

$$P(5, 3) = 5 \times 4 \times 3 = 5!/2! = 60$$

- How many length-4 sequences of distinct digits are there?

$$P(10, 4) = 10 \times 9 \times 8 \times 7 = 10!/6! = 5040$$

- How many four-letter words can be made with distinct letters from the list a, g, m, o, p, r ?

$$P(6, 4) = 6 \times 5 \times 4 \times 3 = 6!/(6-4)! = 360$$

Examples

Counting k -combinations

Examples

- I have a jar with 20 different sweets. Three children come in, and each take one. How many different outcomes are there?

$$P(20, 3) = 20 \times 19 \times 18 = 20!/17! = 6840$$

- I have a jar with 3 different sweets. Three children come in, and each take one. How many different outcomes are there?

$$P(3, 3) = 3 \times 2 \times 1 = 3!/0! = 6$$

A size- k subset is called a **k -combination**

The number of k -combinations of a set of size n is

$$C(n, k) = \frac{n!}{(n-k)!k!}.$$

Proof:

- The number of k -permutations of the set is $P(n, k) = \frac{n!}{n-k}!$
- A k -permutation is an ordering of k distinct elements of the set
- Each size- k subset has $k!$ orderings, so it corresponds to $P(k, k) = k!$ of the k -permutations
- By the division rule, $C(n, k) = \frac{P(n, k)}{P(k, k)} = \frac{n!}{(n-k)!k!}$

- The number of size-2 subsets of $\{1, 2, 3, 4, 5\}$ is

$$C(5, 2) = \frac{5!}{(5-2)!2!} = \frac{5 \times 4 \times 3 \times 2 \times 1}{(3 \times 2 \times 1) \times (2 \times 1)} = \frac{5 \times 4}{2} = 10$$

The ten subsets are $\{1, 2\}$, $\{1, 3\}$, $\{1, 4\}$, $\{1, 5\}$, $\{2, 3\}$, $\{2, 4\}$, $\{2, 5\}$, $\{3, 4\}$, $\{3, 5\}$, and $\{4, 5\}$.

- The number of size-3 subsets of $\{1, 2, 3, 4, 5\}$ is

$$C(5, 3) = \frac{5!}{(5-3)!3!} = 10$$

(The subsets are the complements of the ones above)

Examples	Example	Example continued
<p>■ The number of size 1 subsets of $\{1, 2, 3, 4, 5\}$ is</p> $C(5, 1) = \frac{5!}{(5-1)!1!} = 5,$ <p>which is also the number of size 4 subsets of the set.</p> <p>■ The number of size-0 subsets of $\{1, 2, 3, 4, 5\}$ is</p> $C(5, 0) = \frac{5!}{(5-0)!0!} = 1,$ <p>which is also the number of size-5 subsets $C(5, 5)$.</p>	<p>Twelve people, including Mary and Peter, are candidates to serve on a committee of five. How many different committees are possible? Of these how many</p> <ol style="list-style-type: none"> 1. contain both Mary and Peter? 2. contain neither Mary and Peter? 3. contain either Mary or Peter (but not both)? <p>Solution: There are</p> $C(12, 5) = \frac{12!}{(12-5)!5!} = 792$ <p>possible committees.</p>	<p>(1.) If Mary and Peter are already included, we have to select three more committee members from the remaining ten available people. This can be done in</p> $C(10, 3) = 120$ <p>ways.</p> <p>(2.) If Mary and Peter are excluded we have to select five committee members from the remaining 10 people. This can be done in</p> $C(10, 5) = 252$ <p>ways.</p>
<p>http://www.csc.liv.ac.uk/~konev/COMP109 Part 6. Combinatorics 36 / 44</p>	<p>http://www.csc.liv.ac.uk/~konev/COMP109 Part 6. Combinatorics 37 / 44</p>	<p>http://www.csc.liv.ac.uk/~konev/COMP109 Part 6. Combinatorics 38 / 44</p>

Example continued	Binomial coefficients	The binomial theorem
<p>(3.) The number of committees containing Mary and not Peter is just</p> $C(10, 4)$ <p>The same number of committees contain Peter and exclude Mary. Therefore</p> $2 \times C(10, 4) = 420$ <p>committees contain exactly one of Peter and Mary.</p>	<p>The quantity $C(n, k)$, which gives the number of k-combinations of a set of size n, is called a binomial coefficient.</p> <p>It is also written as</p> $\binom{n}{k} = \frac{n!}{(n-k)!k!}$ <p>(Mathematicians prefer this notation)</p>	<p>For every natural number n,</p> $(a + b)^n = \sum_{k=0}^n \binom{n}{k} a^k b^{n-k}.$ <p>Informally:</p> $(a + b)^n = \underbrace{(a + b) \times (a + b) \times \dots \times (a + b)}_n$ $= \sum_{k=0}^n \sum_{\substack{S \text{ is a} \\ k\text{-combination}}} a^k b^{n-k}$
<p>http://www.csc.liv.ac.uk/~konev/COMP109 Part 6. Combinatorics 39 / 44</p>	<p>http://www.csc.liv.ac.uk/~konev/COMP109 Part 6. Combinatorics 40 / 44</p>	<p>http://www.csc.liv.ac.uk/~konev/COMP109 Part 6. Combinatorics 41 / 44</p>

Pascal's triangle	Binomial coefficient identity	Counting and probabilities																												
<table style="border-collapse: collapse; margin-left: 20px;"> <tr><td style="padding: 2px 10px;">1</td><td style="padding: 2px 10px;"></td><td style="padding: 2px 10px;">$(a + b)^0 = 1$</td></tr> <tr><td style="padding: 2px 10px;">1</td><td style="padding: 2px 10px;">1</td><td style="padding: 2px 10px;">$(a + b)^1 = a + b$</td></tr> <tr><td style="padding: 2px 10px;">1</td><td style="padding: 2px 10px;">2</td><td style="padding: 2px 10px;">1</td><td style="padding: 2px 10px;">$(a + b)^2 = a^2 + 2ab + b^2$</td></tr> <tr><td style="padding: 2px 10px;">1</td><td style="padding: 2px 10px;">3</td><td style="padding: 2px 10px;">3</td><td style="padding: 2px 10px;">1</td><td style="padding: 2px 10px;">$(a + b)^3 = a^3 + 3a^2b + 3ab^2 + b^3$</td></tr> <tr><td style="padding: 2px 10px;">1</td><td style="padding: 2px 10px;">4</td><td style="padding: 2px 10px;">6</td><td style="padding: 2px 10px;">4</td><td style="padding: 2px 10px;">1</td><td style="padding: 2px 10px;">$(a + b)^4 = a^4 + 4a^3b + 6a^2b^2 + 4ab^3 + b^4$</td></tr> <tr><td style="padding: 2px 10px;">1</td><td style="padding: 2px 10px;">5</td><td style="padding: 2px 10px;">10</td><td style="padding: 2px 10px;">10</td><td style="padding: 2px 10px;">5</td><td style="padding: 2px 10px;">1</td><td style="padding: 2px 10px;">$(a + b)^5 = a^5 + 5a^4b + 10a^3b^2 + 10a^2b^3 + 5ab^4 + b^5$</td></tr> </table>	1		$(a + b)^0 = 1$	1	1	$(a + b)^1 = a + b$	1	2	1	$(a + b)^2 = a^2 + 2ab + b^2$	1	3	3	1	$(a + b)^3 = a^3 + 3a^2b + 3ab^2 + b^3$	1	4	6	4	1	$(a + b)^4 = a^4 + 4a^3b + 6a^2b^2 + 4ab^3 + b^4$	1	5	10	10	5	1	$(a + b)^5 = a^5 + 5a^4b + 10a^3b^2 + 10a^2b^3 + 5ab^4 + b^5$	$\binom{n+1}{r+1} = \binom{n}{r} + \binom{n}{r+1}.$ <p>Proof:</p> <ul style="list-style-type: none"> ■ $\binom{n+1}{r+1}$ is the number of ways to choose a subset of size $r + 1$ from a set of size $n + 1$. ■ Suppose the set is $\{x_1, x_2, \dots, x_{n+1}\}$. ■ How many subsets include x_{n+1}? We just choose a size-r subset from $\{x_1, x_2, \dots, x_n\}$, so there are $\binom{n}{r}$ ways to do it. ■ How many subsets exclude x_{n+1}? We just have to choose a subset of size $r + 1$ from $\{x_1, x_2, \dots, x_n\}$, so there are $\binom{n}{r+1}$ ways to do it. ■ These outcomes are disjoint, so the total number of subsets is the sum of $\binom{n}{r}$ and $\binom{n}{r+1}$. 	<p>Counting also helps us to answer questions like</p> <ul style="list-style-type: none"> ■ What are the odds of winning the National Lottery? ■ What payoffs should you expect? ■ What is more likely in poker, full house or 4-of-a-kind?
1		$(a + b)^0 = 1$																												
1	1	$(a + b)^1 = a + b$																												
1	2	1	$(a + b)^2 = a^2 + 2ab + b^2$																											
1	3	3	1	$(a + b)^3 = a^3 + 3a^2b + 3ab^2 + b^3$																										
1	4	6	4	1	$(a + b)^4 = a^4 + 4a^3b + 6a^2b^2 + 4ab^3 + b^4$																									
1	5	10	10	5	1	$(a + b)^5 = a^5 + 5a^4b + 10a^3b^2 + 10a^2b^3 + 5ab^4 + b^5$																								
<p>http://www.csc.liv.ac.uk/~konev/COMP109 Part 6. Combinatorics 42 / 44</p>	<p>http://www.csc.liv.ac.uk/~konev/COMP109 Part 6. Combinatorics 43 / 44</p>	<p>http://www.csc.liv.ac.uk/~konev/COMP109 Part 6. Combinatorics 44 / 44</p>																												