

I-Maps: Graphs and Distributions

Dr. Xiaowei Huang

<https://cgi.csc.liv.ac.uk/~xiaowei/>

- Briefing about Assignment 2

Up to now,

- Overview of Machine Learning
- Traditional Machine Learning Algorithms
- Deep learning
- Probabilistic Graphical Models
 - Introduction

Topics

- Recap: Conditional Independence
- Markov Assumption and Definition of I-Maps
- I-Map to Factorization
- Factorization to I-Map
- Perfect Map

Graphs and Distributions

- Relating two concepts:
 - Conditional Independencies in distributions
 - Conditional Independencies in graphs
- I-Map is a relationship between the two

Recap: Conditional Independence

Recap: Conditional Independence

- Two variables X and Y are **conditionally independent** given Z if
 - $P(X = x|Y = y, Z = z) = P(X = x|Z = z)$ for all values x, y, z
 - That is, learning the values of Y does not change prediction of X once we know the value of Z
 - notation: $(X \perp Y | Z)$

Recap: Conditional Independence

- X, Y independent $X \perp Y$ or $X \perp Y | \emptyset$

if and only if:

$$\forall x, y : P(x, y) = P(x)P(y)$$

- X and Y are conditionally independent given Z: $X \perp Y | Z$

if and only if:

$$\forall x, y, z : P(x, y | z) = P(x | z)P(y | z)$$

Independencies in a Distribution

- Let P be a distribution over X
- Define $I(P)$ to be the set of conditional independence assertions of the form $(X \perp Y | Z)$ that hold in P
- Example:

X	Y	P(X,Y)
x^0	y^0	0.08
x^0	y^1	0.32
x^1	y^0	0.12
x^1	y^1	0.48

X and Y are independent in P , e.g.,

$$P(x^1) = 0.48 + 0.12 = 0.6$$

$$P(y^1) = 0.32 + 0.48 = 0.8$$

$$P(x^1, y^1) = 0.48 = 0.6 \times 0.8$$

Thus $(X \perp Y | \phi) \in I(P)$

Independencies in a Distribution

- Let P be a distribution over X
- Define $I(P)$ to be the set of conditional independence assertions of the form $(X \perp Y | Z)$ that hold in P
- Example:

X	Y	P(X,Y)
x^0	y^0	0.08
x^0	y^1	0.32
x^1	y^0	0.12
x^1	y^1	0.48

X and Y are independent in P , e.g.,

$$P(x^1) = 0.48 + 0.12 = 0.6$$

$$P(y^1) = 0.32 + 0.48 = 0.8$$

$$P(x^1, y^1) = 0.48 = 0.6 \times 0.8$$

Thus $(X \perp Y | \phi) \in I(P)$

How about this distribution?


X	Y	P(X,Y)
x^0	y^0	0.10
x^0	y^1	0.16
x^1	y^0	0.64
x^1	y^1	0.10

Markov Assumption and Definition of I-Map


Markov Assumption

- We now make this independence assumption more precise for **directed acyclic graphs** (DAGs)
- Each random variable X , is independent of its non-descendants, given its parents $Pa(X)$
- Formally,

$$(X \perp NonDesc(X) | pa(X))$$



Can we read off the independencies from a graph?



Independencies in a Graph


- Graph G with CPDs is equivalent to a set of independence assertions

$$P(D, I, G, S, L) = P(D)P(I)P(G | D, I)P(S | I)P(L | G)$$

- Local Conditional Independence Assertions (starting from leaf nodes):

$I(G) = \{(L \perp I, D, S | G),$ L is conditionally independent of all other nodes given parent G
 $(S \perp D, G, L | I),$ S is conditionally independent of all other nodes given parent I
 $(G \perp S | D, I),$ Even given parents, G is NOT independent of descendant L
 $(I \perp D | \phi),$ Nodes with no parents are marginally independent
 $(D \perp I, S | \phi)\}$ D is independent of non-descendants I and S


- Parents of a variable shield it from probabilistic influence
 - Once value of parents known, no influence of ancestors
- Information about descendants can change beliefs about a node




Definition of I-MAP

- Let G be a graph associated with a set of independencies $I(G)$
- Let P be a probability distribution with a set of independencies $I(P)$
- Then G is an **I-Map** of P if $I(G) \subseteq I(P)$
 - Intuitively, a DAG G is an **I-Map** of a distribution P if all Markov assumptions implied by G are satisfied by P
- From direction of inclusion
 - distribution can have more independencies than the graph
 - Graph does not mislead in independencies existing in P
 - Any independence that G asserts must also hold in P


Example of I-MAP



G_0 encodes
 $X \perp Y$ or
 $I(G_0) = \{X \perp Y\}$




G_1 encodes no
Independence, or
 $I(G_1) = \emptyset$




G_2 encodes no
Independence, or
 $I(G_2) = \emptyset$


Example of I-MAP



G_0 encodes
 $X \perp Y$ or
 $I(G_0) = \{X \perp Y\}$



G_1 encodes no
Independence, or
 $I(G_1) = \emptyset$



G_2 encodes no
Independence, or
 $I(G_2) = \emptyset$


X	Y	$P(X,Y)$
x^0	y^0	0.08
x^0	y^1	0.32
x^1	y^0	0.12
x^1	y^1	0.48

X and Y are independent
in P , e.g.,


G_0 is an I-map of P
 G_1 is an I-map of P
 G_2 is an I-map of P

If G is an I-map of P then it captures **some** of the independences, not all


Example of I-MAP



G_0 encodes
 $X \perp Y$ or
 $I(G_0) = \{X \perp Y\}$



G_1 encodes no
Independence, or
 $I(G_1) = \emptyset$



G_2 encodes no
Independence, or
 $I(G_2) = \emptyset$

X	Y	$P(X,Y)$
x^0	y^0	0.4
x^0	y^1	0.3
x^1	y^0	0.2
x^1	y^1	0.1

X and Y are not
independent in P
Thus $(X \perp Y) \notin I(P)$

G_0 is not an I-map of P
 G_1 is an I-map of P
 G_2 is an I-map of P

If G is an I-map of P then it captures **some** of the independences, not all

Exercise

- Please draw an I-Map for each of the following distributions:

x	y	P(x,y)
0	0	0.25
0	1	0.25
1	0	0.25
1	1	0.25

x	y	P(x,y)
0	0	0.2
0	1	0.3
1	0	0.4
1	1	0.1

I-map to Factorization

What is factorization?

- **factorization** or **factoring** consists of writing a number or another mathematical object as a product of several *factors*, usually smaller or simpler objects of the same kind
- In our context, for example:

$$P(D, I, G, S, L) = P(D)P(I)P(G|D, I)P(S|I)P(L|G)$$

or

$$P(I, D, G, L, S) = P(I)P(D|I)P(G|I, D)P(L|I, D, G)P(S|I, D, G, L)$$


I-map to Factorization

- Consider Joint distribution $P(I, D, G, L, S)$

- From chain rule of probability

$$P(I, D, G, L, S) = P(I)P(D|I)P(G|I, D)P(L|I, D, G)P(S|I, D, G, L)$$

- Relies on no assumptions, also not very helpful
 - Last factor requires evaluation of 24 conditional probabilities



Factorization Theorem


- **Thm:** if G is an I-Map of P , then

$$P(X_1, \dots, X_n) = \prod_i P(X_i \mid Pa(X_i))$$

I-map to Factorization

- Start from $P(I, D, G, L, S) = P(I)P(D|I)P(G|I, D)P(L|I, D, G)P(S|I, D, G, L)$
- Assume G is an I-map
 - Apply conditional independence assumptions induced from the graph

- *we have* $(D \perp I) \in I(G) \subseteq I(P)$
- *and thus* $P(D|I) = P(D)$



- *Therefore, we have*

$$P(I, D, G, L, S) = P(I)P(D)P(G|I, D)P(L|I, D, G)P(S|I, D, G, L)$$

We can go from graphs to factorization of P

I-map to Factorization


- Start from $P(I, D, G, L, S) = P(I)P(D|I)P(G|I, D)P(L|I, D, G)P(S|I, D, G, L)$

- We have $(L \perp I, D | G) \in I(G) \subseteq I(P)$

- And thus $P(L|I, D, G) = P(L|G)$

- Therefore, we have

$$P(I, D, G, L, S) = P(I)P(D)P(G|I, D)P(L|G)P(S|I, D, G, L)$$



We can go from graphs to factorization of P

I-map to Factorization


- Start from $P(I, D, G, L, S) = P(I)P(D)P(G|I, D)P(L|G)P(S|I, D, G, L)$

- We have $(S \perp D, G, L | I) \in I(G) \subseteq I(P)$

- And thus $P(S|I, D, G, L) = P(S|I)$

- Therefore, we have


$$P(I, D, G, L, S) = P(I)P(D)P(G|I, D)P(L|G)P(S|I)$$



We can go from graphs to factorization of P

Exercise

- Please give the factorization of the distribution P according to the I-Map shown in the figure.



Factorization to l-map

Factorization to I-map


- We can also show the opposite

Thm

$$P(X_1, \dots, X_n) = \prod_i P(X_i \mid Pa_i) \Rightarrow \mathbf{G} \text{ is an I-Map of } P$$

Proof (Outline) Assume we have graph G , we need to prove independence

$$\begin{aligned} P(Z \mid X, Y) &= \frac{P(X, Y, Z)}{P(X, Y)} = \frac{P(X)P(Y \mid X)P(Z \mid X)}{P(X)P(Y \mid X)} \\ &= P(Z \mid X) \end{aligned}$$




Factorization to I-map

- We have seen that we can go from the independences encoded in G , i.e., $I(G)$, to Factorization of P
- Conversely, We have factorization and associated G . *We can show* that conditional independences from $I(G)$ are all in $I(P)$.

Example that independences in G hold in P

- P is defined by set of CPDs
- Starting from factorization and its associated graph

$$P(D, I, G, S, L) = P(I)P(D)P(G|I, D)P(L|G)P(S|I)$$



- Consider independences for S in G , i.e.,

$$P(S \perp D, G, L | I)$$

- We need to show that $P(S|I, D, G, L) = P(S|I)$

Exercise?

Perfect Map

Perfect Map

- I-map
 - All independencies in $I(G)$ present in $I(P)$
 - Trivial case: all nodes interconnected
- D-Map
 - All independencies in $I(P)$ present in $I(G)$
 - Trivial case: all nodes disconnected
- Perfect map
 - Both an I-map and a D -map
 - Interestingly not all distributions P over a given set of variables can be represented as a perfect map
 - Venn Diagram where D is set of distributions that can be represented as a perfect map

